



## Characterizing *Yarrowia lipolytica* as a cell factory for sustainable bioprocesses

Lubuta, Patrice; Workman, Mhairi

*Publication date:*  
2016

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Lubuta, P., & Workman, M. (2016). *Characterizing Yarrowia lipolytica as a cell factory for sustainable bioprocesses*. Poster session presented at 12th International Conference on Renewable Resources and Biorefineries, Ghent, Belgium.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.





# Characterizing *Yarrowia lipolytica* as a cell factory for sustainable bioprocesses

Patrice Lubuta and Mhairi Workman

Fungal Physiology and Biotechnology Group  
Department of Systems Biology, Technical University of Denmark,  
Building 223, 2800 Kgs. Lyngby, Denmark

## Background

The utilization of an increasing amount of agro-waste and renewable low-cost substrates will be an ongoing challenge for the bio-based economy. Mobilization of these feedstocks is necessary for realization of biosustainability on an industrial scale. Cell factories traditionally applied in biotechnology are using a limited range of substrates (mainly glucose). However, a wider diversity in substrate range is highly desirable in developing biorefinery scenarios where seasonal variation in availability and composition of biomass feed-stocks typically appear.

## Aim of the study

This study evaluates the applicability of the non-conventional yeast *Y. lipolytica* as a cell factory for the conversion of **glycerol** and the lignocellulose hydrolysate sugars **glucose**, **xylose** and **arabinose**. A quantitative physiological investigation was carried out in order to assess the cellular performance of three different *Y. lipolytica* wild type strains (**IBT 446**, **W29** and **H222**) in single and mixed carbon source cultivations.

## Benchmarking strains on single carbon substrate

### Glucose vs Glycerol

	IBT 446		W29		H222	
	Glucose <sup>1</sup>	Glycerol <sup>1</sup>	Glucose	Glycerol	Glucose	Glycerol
<b>Growth rate</b>						
$\mu_{max}$ (h <sup>-1</sup> )	0,24 ± 0,01	0,30 ± 0,01	0,28 ± 0,02	0,31 ± 0,02	0,28 ± 0,01	0,35 ± 0,01
<b>Yield coefficients</b>						
Y <sub>sx</sub> (cmol cmol <sup>-1</sup> )	0,69 ± 0,03	0,61 ± 0,01	0,67 ± 0,03	0,73 ± 0,01	0,64 ± 0,02	0,72 ± 0,01
Y <sub>sc</sub> (cmol cmol <sup>-1</sup> )	0,30 ± 0,01	0,23 ± 0,00	0,34 ± 0,02	0,30 ± 0,01	0,35 ± 0,004	0,21 ± 0,01
Y <sub>sm</sub> (cmol cmol <sup>-1</sup> )	N/A*	0,17 ± 0,00	N/A*	0,015 ± 0,01	N/A*	N/A**
<b>Total</b>	<b>0,99 ± 0,03</b>	<b>1,01 ± 0,01</b>	<b>1,02 ± 0,05</b>	<b>1,04 ± 0,01</b>	<b>0,98 ± 0,01</b>	<b>0,94 ± 0,01</b>

<sup>1</sup>Workman 2013, N/A: Not applicable; \*below detection limit. \*\*Carbon source not depleted at last sample point.

### Xylose and Arabinose

Growth experiments in shake flasks and defined minimal media with 0.65 cmol L<sup>-1</sup> (≈ 20 g L<sup>-1</sup>) of the carbon source and cultivation time of 140 h: **None of the strains showed growth on xylose or arabinose as the sole carbon source (verified by OD and HPLC measurements).**

**Set up:** Bioreactor batch cultivation, minimal media with 0,163 cmol L<sup>-1</sup> (≈ 5 g L<sup>-1</sup>) of each carbon source.

### IBT 446 - mixed substrates



#### Results:

Uptake cascade: 1. glycerol, 2. glucose, 3. xylose, 4. arabinose

Xylitol production and re-utilization

Growth on glycerol and glucose (primary carbon sources)

Maintenance of biomass after depletion of primary carbon sources



### W29 - mixed substrates

#### Results:

Uptake cascade: 1. glycerol, 2. glucose, 3. xylose (half consumed)


Co-consumption glycerol/glucose

Arabinose not consumed

Xylitol production but no re-utilization

Growth on glycerol and glucose (primary carbon sources)

Drop of biomass after depletion of primary carbon sources



### W29 - mixed substrates

#### Results:

Uptake cascade: 1. glycerol, 2. glucose, 3. xylose (half consumed)

Arabinose not consumed

Xylitol production but no re-utilization

Growth on glycerol and glucose (primary carbon sources)

Drop of biomass after depletion of primary carbon sources

## Summary and Conclusions

With current global focus on renewable and sustainable technologies, there is a strong need to develop innovative solutions in industrial biotechnology. The non-conventional yeast *Yarrowia lipolytica* has considerable potential as a versatile cell factory, especially when compared to the typically applied organisms such as *Saccharomyces cerevisiae* which utilizes glycerol very slowly and requires genetic engineering for conversion of lignocellulosic sugar monomers xylose and arabinose. In this study we could show that *Y. lipolytica* IBT 446 is versatile and interesting as a biorefinery cell factory, as it is natural capable to utilize sugars typically found in plant hydrolysates (glucose, xylose, arabinose) together with glycerol in mixed carbon cultivations. Furthermore the strains W29 and H222 are naturally capable of conversion of glycerol, glucose and xylose.